arXiv:1611.00662v1 [math.RT] 2 Nov 2016

CAPELLI ELEMENTS OF THE GROUP ALGEBRA

NAOYA YAMAGUCHI

ABSTRACT. Inspired by the Capelli identities for group determinants obtained by Tôru Umeda, we give a basis of the center of the group algebra of any finite group by using Capelli identities for irreducible representations.

The Capelli identities for irreducible representations are modifications of the Capelli identity. These identities lead to Capelli elements of the group algebra. These elements construct a basis of the center of the group algebra.

1. INTRODUCTION

The Capelli identity is analogous to the product formula for the determinant in the Weyl algebra. The identity leads to the Capelli element. It is known that the Capelli element is a central element in the universal enveloping algebra of $\mathfrak{gl}_n$.

On the other hand, An Huang gave Capelli-type identities associated with the quaternions and the octonions [3]. Inspired by his results, Tôru Umeda gave Capelli identities for group determinants [7]. There are Capelli identities for irreducible representations in the background of the Capelli identities for group determinants.

In this paper, we give a basis of the center of the group algebra of any finite group by using Capelli identities for irreducible representations. These identities lead to Capelli elements of the group algebra. These elements construct a basis.

First, we explain our motivation.

1.1. Motivation. Let G be a finite group, $\widehat{G}$ a complete set of irreducible representations of G over $\mathbb{C}$, $\mathbb{C}G = \left\{\sum_{g\in G} x_g g \mid x_g \in \mathbb{C}\right\}$ the group algebra, and $Z(\mathbb{C}G)$ the center of $\mathbb{C}G$. The following theorem is easily proven from Schur's orthogonal relations.

Theorem 1. *Let χ_φ be the character of $\varphi \in \widehat{G}$. The set*

$$\left\{ \mathrm{Tr}\left(\sum_{g\in G} \varphi(g)g\right) \mid \varphi \in \widehat{G} \right\} = \left\{ \sum_{g\in G} \chi_\varphi(g)g \mid \varphi \in \widehat{G} \right\}$$

is a basis of $Z(\mathbb{C}G)$ where we omit the numbering of the element of the basis.

At this point, we have a simple question. Is the set $\left\{\det\left(\sum_{g\in G}\varphi(g)g\right) \mid \varphi \in \widehat{G}\right\}$ a basis of $Z(\mathbb{C}G)$? Our main result gives an answer.

Date: November 3, 2016.

2010 *Mathematics Subject Classification.* Primary 20C05; Secondary 15A15.

Key words and phrases. Capelli identity; Capelli element; group algebra.

1.2. Main result. Let z be a complex variable, $|G|$ the order of G, $m = \deg\varphi$, $\alpha = \frac{|G|}{m}$, $u_i(z) = \alpha(m-i) - z$, $u^{(i)}(z) = u_m(z)u_{m-1}(z)\cdots u_{m-i+1}(z)$, det the column determinant, and the Capelli element for φ of the group algebra

$$\overline{C}^{\varphi}(z) = \det\left(\sum_{g\in G}\varphi(g)g + \alpha\left(\begin{bmatrix} m-1 & & & \\ & m-2 & & \\ & & \ddots & \\ & & & 0 \end{bmatrix}\right) - zI_m\right) \in \mathbb{C}[z]\otimes\mathbb{C}G.$$

Then we can prove the following relation.

Theorem 2. *We have*

$$\overline{C}^{\varphi}(z) = u^{(m)}(z) + \mathrm{Tr}\left(\sum_{g\in G}\varphi(g)g\right)u^{(m-1)}(z)$$

The above relation leads to the following corollary.

Corollary 3. *Suppose $k_\varphi \in \mathbb{C}$ such that $u^{(m-1)}(k_\varphi) \neq 0$. Then,*

$$\left\{\overline{C}^{\varphi}(k_\varphi) \mid \varphi \in \widehat{G}\right\}$$

is a basis of $Z(\mathbb{C}G)$.

This is our answer. We provide some sections for the details.

2. Capelli identity and Capelli element

Here, we review the Capelli identity and the Capelli element.

2.1. Column determinant. First, we explain the column determinant. Let R be an associative algebra.

Definition 4 (Column determinant)**.** *Let $A = (a_{ij})_{1\leq i,j\leq m} \in \mathrm{Mat}(m,R)$. We define the column determinant of A by*

$$\det A = \sum_{\sigma\in S_m}\mathrm{sgn}(\sigma)a_{\sigma(1)1}a_{\sigma(2)2}\cdots a_{\sigma(m)m}.$$

Hence, we have $\det\begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - cb$.

2.2. Weyl algebra. The Capelli identity is analogous to the product formula for the determinant in the Weyl algebra. Next, we explain the Weyl algebra $\mathbb{C}[x_{ij}, \partial_{kl} \mid 1 \leq i,j,k,l \leq m]$.

Let x_{ij} $(1 \leq i,j \leq m)$ be variables and $\partial_{ij} = \frac{\partial}{\partial x_{ij}}$ $(1 \leq i,j \leq m)$ partial differential operators. We assume that these variables and operators are related as follows.

For all $1 \leq i,j,k,l \leq m$, we have

$$[x_{ij}, x_{kl}] = 0, \quad [\partial_{ij}, \partial_{kl}] = 0, \quad [\partial_{ij}, x_{kl}] = \alpha\delta_{ik}\delta_{jl}$$

where δ is the Kronecker delta. Usually, we take $\alpha = 1$. Here, we will not assume that $\alpha = 1$. The Weyl algebra is generated by these variables and operators.

2.3. Capelli identity. Next, we explain the Capelli identity. Let

$$X = (x_{ij})_{1\leq i\leq m, 1\leq j\leq m}, \qquad \partial = (\partial_{ij})_{1\leq i\leq m, 1\leq j\leq m},$$
$$\Pi = {}^tX\partial, \qquad \natural_m = \mathrm{diag}(m-1, m-2, \ldots, 0)$$

The Capelli identity is as follows.

Theorem 5 (Capelli identity)**.** *We have*

$$\det(\Pi + \alpha\natural_m) = \det X \det \partial.$$

Example 6. *Let $m = 2$ and $\alpha = 1$. We have*

$$\det\begin{bmatrix} x_{11}\partial_{11} + x_{21}\partial_{21} + 1 & x_{11}\partial_{12} + x_{21}\partial_{22} \\ x_{12}\partial_{11} + x_{22}\partial_{21} & x_{21}\partial_{12} + x_{22}\partial_{12} \end{bmatrix} = \det\begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{bmatrix} \det\begin{bmatrix} \partial_{11} & \partial_{12} \\ \partial_{21} & \partial_{22} \end{bmatrix}$$

2.4. Capelli element. The Capelli element is a charactrestic polynomial of Π. Let z be a variable.

Definition 7 (Capelli element)**.** *We define the Capelli element $C(z)$ by*

$$C(z) = \det(\Pi + \alpha\natural_m - zI_m).$$

The Capelli identity is conjugation invariant.

Theorem 8. *For all $P \in GL(m, \mathbb{C})$, we have*

$$\det\left(P\Pi P^{-1} + \alpha\natural_m - zI_m\right) = C(z).$$

The following theorem plays an important role in what follows.

Theorem 9. *For all $1 \leq i, j \leq m$, we have*

$$[\Pi_{ij}, C(z)] = 0.$$

Theorem 8 and 9 are obtained from only the following relations.
For all $1 \leq i, j, k, l \leq m$,

$$[\Pi_{ij}, \Pi_{kl}] = \alpha(\delta_{jk}\Pi_{il} - \delta_{il}\Pi_{kj}).$$

3. Capelli identity for irreducible representations

Here, we explain the Capelli identities for irreducible representations.

Let G be a finite group, x_g $(g \in G)$ variable and $\partial_g = \frac{\partial}{\partial x_g}$ $(g \in G)$ partial differential operator. We assume that the following relations hold.

For all $g, h \in G$,

$$[x_g, x_h] = 0, \quad [\partial_g, \partial_h] = 0, \quad [\partial_g, x_h] = \delta_{gh}.$$

Then, we have the Weyl algebra $\mathbb{C}[x_g, \partial_h]$. Next, we construct Weyl subalgebras of the Weyl algebra by using irreducible unitary representations of G.

Let $|G|$ be the cardinality of the set G (that is, $|G|$ is the order of the group G), φ a unitary matrix form of an irreducible representation of G,

$$\alpha = \frac{|G|}{m}, \quad X^{\varphi} = \sum_{g\in G} \overline{\varphi(g)} x_g, \quad \partial^{\varphi} = \sum_{g\in G} \varphi(g)\partial_g, \text{and} \quad \Pi^{\varphi} = {}^tX^{\varphi}\partial^{\varphi}$$

where $\overline{\varphi(g)}$ is the complex conjugate matrix of $\varphi(g)$. Then, we have the following relations.

For all $1 \leq i, j, k, l \leq m$,

$$[X_{ij}^{\varphi}, X_{kl}^{\varphi}] = 0, \quad [\partial_{ij}^{\varphi}, \partial_{kl}^{\varphi}] = 0, \quad [\partial_{ij}^{\varphi}, X_{kl}^{\varphi}] = \alpha \delta_{ik} \delta_{jl}.$$

This leads us to the following identity.

Theorem 10 (Capelli identity for irreducible representations)**.** *We have*

$$\det(\Pi^{\varphi} + \alpha \natural_m) = \det X^{\varphi} \det \partial^{\varphi}.$$

Let $C^{\varphi}(z) = \det(\Pi^{\varphi} + \alpha \natural_m - zI_m)$ be the Capelli element. From Theorem 8, the Capelli element is invariant under a change of a matrix form of the irreducible representation. This enables us to redefine the Capelli element as follows.

Definition 11 (Capelli element for irreducible representations)**.** *Let $\varphi \in \widehat{G}$ and $m = \deg \varphi$. We define $C^{\varphi}(z)$ by*

$$C^{\varphi}(z) = \det(\Pi^{\varphi} + \alpha \natural_m - zI_m)$$

We call $C^{\varphi}(z)$ the Capelli element for φ.

4. Capelli element of the group algebra

Let $\mathbb{C}G = \left\{ \sum_{g \in G} x_g g \mid x_g \in \mathbb{C} \right\}$ the group algebra of G, $\widetilde{G}$ a complete set of irreducible unitary matrix representations of G, $\varphi \in \widetilde{G}$, and

$$E^{\varphi} = \sum_{g \in G} \varphi(g) g \in \mathrm{Mat}(\deg \varphi, \mathbb{C}G).$$

From Schur's orthogonal relations, we have the following lemmas.

Lemma 12. *$\{E_{ij}^{\varphi} \mid 1 \leq i, j \leq \deg \varphi, \varphi \in \widetilde{G}\}$ is a basis of $\mathbb{C}G$.*

Lemma 13. *Let $\varphi, \psi \in \widetilde{G}$, where φ is not equivalent to ψ. For all $1 \leq i, j \leq \deg \varphi$ and $1 \leq s, t \leq \deg \psi$, we have*

$$E_{ij}^{\varphi} E_{kl}^{\varphi} = \alpha_{\deg \varphi} \delta_{jk} E_{il}, \quad E_{ij}^{\varphi} E_{st}^{\psi} = 0.$$

In particular, we have

$$[E_{ij}^{\varphi}, E_{kl}^{\varphi}] = \alpha_{\deg \varphi}(\delta_{jk} E_{il} - \delta_{il} E_{kj}), \tag{1}$$

$$[E_{ij}^{\varphi}, E_{kl}^{\psi}] = 0. \tag{2}$$

Let

$$\overline{C}^{\varphi}(z) = \det(E^{\varphi} + \alpha_m \natural_m - zI_m) \in \mathbb{C}[z] \otimes \mathbb{C}G.$$

Recall that Theorem 8 and Theorem 9 are obtained from only the relations $[\Pi_{ij}, \Pi_{kl}] = \alpha(\delta_{jk}\Pi_{il} - \delta_{il}\Pi_{kj})$. Hence, $\overline{C}^{\varphi}(z)$ is conjugation invariant from the relations (1), and we have

$$[E_{ij}^{\varphi}, \overline{C}^{\varphi}(z)] = 0 \tag{3}$$

for any $1 \leq i, j \leq \deg \varphi$.

Using the above conjugation invariance, we redefine $\overline{C}^{\varphi}(z)$.

Definition 14 (Capelli element of the group algebra)**.** *Let $\varphi \in \widehat{G}$. We define the Capelli element for φ of the group algebra by*

$$\overline{C}^{\varphi}(z) = \det(E^{\varphi} + \alpha_m \natural_m - zI_m).$$

From Lemma 12 and relations (1) and (2), we can prove the following Lemma.

Lemma 15. *For all $\varphi \in \widehat{G}$, $\overline{C}^{\varphi}(z) \in Z(\mathbb{C}G)$. That is, $\overline{C}^{\varphi}(z)$ is a central element of the group algebra.*

Let $u_i(z) = \alpha_m(m-i) - z$, $u^{(i)}(z) = u_m(z)u_{m-1}(z)\cdots u_{m-i+1}(z)$, $E_{ij}^{\varphi}(u_i(z)) = E_{ij}^{\varphi} + \delta_{ij}u_i(z)$, and $[m] = \{1, 2, \ldots, m\}$. The following is the main theorem.

Theorem 16. *We have*

$$\overline{C}^{\varphi}(z) = u^{(m)}(z) + \mathrm{Tr}(E^{\varphi})u^{(m-1)}(z)$$

Proof. From the definition of $\overline{C}^{\varphi}(z)$, we have

$$\begin{aligned}
\overline{C}^{\varphi}(z) &= \sum_{\sigma\in S_m} \mathrm{sgn}(\sigma) E^{\varphi}_{\sigma(1)1}(u_1(z)) E^{\varphi}_{\sigma(2)2}(u_2(z)) \cdots E^{\varphi}_{\sigma(m)m}(u_m(z)) \\
&= \sum_{\sigma\in S_m} \mathrm{sgn}(\sigma)(E^{\varphi}_{\sigma(1)1} + \delta_{\sigma(1)1}u_1(z)) \cdots (E^{\varphi}_{\sigma(m)m} + \delta_{\sigma(m)m}u_m(z)) \\
&= \sum_{\sigma\in S_m} \mathrm{sgn}(\sigma) \sum_{T\subset[m]} \prod_{t\in T} E^{\varphi}_{\sigma(t)t} \prod_{s\in[m]\setminus T} \delta_{\sigma(s)s}u_s(z).
\end{aligned}$$

Let $T = \{t_1 < t_2 < \ldots < t_{|T|}\}$ and $[m]\setminus T = \{s_1, s_2, \ldots, s_{m-|T|}\}$. From Lemma 13, we have

$$\begin{aligned}
&\sum_{\sigma\in S_m} \mathrm{sgn}(\sigma) \sum_{T\subset[m]} \prod_{t\in T} E^{\varphi}_{\sigma(t)t} \prod_{s\in[m]\setminus T} \delta_{\sigma(s)s}u_s(z) \\
&= \alpha_m^{|T|-1} \sum_{\sigma\in S_m} \mathrm{sgn}(\sigma)\delta_{\sigma(t_2)t_1}\delta_{\sigma(t_3)t_2}\cdots\delta_{\sigma(t_{|T|})t_{|T|-1}} E^{\varphi}_{\sigma(t_1)t_{|T|}} \prod_{s\in[m]\setminus T} \delta_{\sigma(s)s}u_s(z) \\
&= (-\alpha_m)^{|T|-1} E^{\varphi}_{t_{|T|}t_{|T|}} \prod_{s\in[m]\setminus T} u_s(z).
\end{aligned}$$

Therefore, there exists $a_i \in \mathbb{C}[z]$ $(1 \le i \le m)$ such that

$$\overline{C}^{\varphi}(z) = \sum_{i=1}^{m} a_i E^{\varphi}_{ii} + u^{(m)}(z).$$

We show that $a_p = a_q$ for all $p, q \in [m]$. From Lemma 13, we have

$$\begin{aligned}
E^{\varphi}_{pq}\overline{C}^{\varphi}(z) &= E^{\varphi}_{pq}\left(\sum_{i=1}^{m} a_i E^{\varphi}_{ii} + u^{(m)}(z)\right) \\
&= \sum_{i=1}^{m} a_i\delta_{iq}\alpha_m E^{\varphi}_{pi} + u^{(m)}(z)E^{\varphi}_{pq} \\
&= a_q\alpha_m E^{\varphi}_{pq} + u^{(m)}(z)E^{\varphi}_{pq}, \\
\overline{C}^{\varphi}(z)E^{\varphi}_{pq} &= \left(\sum_{i=1}^{m} a_i E^{\varphi}_{ii} + u^{(m)}(z)\right) E^{\varphi}_{pq} \\
&= \sum_{i=1}^{m} a_i\delta_{ip}\alpha_m E^{\varphi}_{iq} + u^{(m)}(z)E^{\varphi}_{pq} \\
&= a_p\alpha_m E^{\varphi}_{pq} + u^{(m)}(z)E^{\varphi}_{pq}.
\end{aligned}$$

From Lemma 15, we have $a_p = a_q$ for all $p, q \in [m]$. We calculate a_1. From $\overline{C}^{\varphi}(z) = \sum_{T\subset[m]}(-\alpha_m)^{|T|-1}E^{\varphi}_{t_{|T|}t_{|T|}}\prod_{s\in[m]\setminus T}u_s(z)$, we have

$$\begin{aligned} a_1E^{\varphi}_{11} &= (-\alpha_m)^{\{1\}-1}E^{\varphi}_{11}\prod_{s\in[m]\setminus\{1\}}u_s(z) \\ &= u^{(m-1)}(z)E^{\varphi}_{11}. \end{aligned}$$

This completes the proof. □

In addition, we have the following corollary.

Corollary 17. *Suppose $k_\varphi \in \mathbb{C}$ such that $u^{(m-1)}(k_\varphi) \neq 0$. Then,*

$$\left\{\overline{C}^{\varphi}(k_\varphi) \mid \varphi \in \widehat{G}\right\}$$

is a basis of $Z(\mathbb{C}G)$.

5. Relationship between column, row and double determinant

In this last section, we explain the relationship between column, row, and double determinants. The row and double determinants are as follows.

Definition 18 (Row determinant)**.** *Let $A = (a_{ij})_{1\leq i,j\leq m} \in \mathrm{Mat}(m, R)$. We define the row determinant of A is defined as*

$$\mathrm{rdet}A = \sum_{\sigma\in S_m}\mathrm{sgn}(\sigma)a_{1\sigma(1)}a_{2\sigma(2)}\cdots a_{m\sigma(m)}.$$

Definition 19 (Double determinant)**.** *Let $A = (a_{ij})_{1\leq i,j\leq m} \in \mathrm{Mat}(m, R)$. The double determinant of A is defined as*

$$\mathrm{Det}A = \frac{1}{m!}\sum_{\sigma,\tau\in S_m}\mathrm{sgn}(\sigma\tau)a_{\sigma(1)\tau(1)}a_{\sigma(2)\tau(2)}\cdots a_{\sigma(m)\tau(m)}.$$

Reference [5] describes that the relationship between column, row, and double determinants. Let

$$\natural^* = \begin{bmatrix} 0 & & & \\ & 1 & & \\ & & \ddots & \\ & & & m-1 \end{bmatrix}, \quad \natural_\sigma = \begin{bmatrix} \sigma(m) & & & \\ & \sigma(m-1) & & \\ & & \ddots & \\ & & & \sigma(1) \end{bmatrix} \quad (\sigma \in S_m),$$

and $E \in \mathrm{Mat}(m, R)$, where we assume that $[E_{ij}, E_{kl}] = \delta_{jk}E_{il} - \delta_{il}E_{kj}$ for all $1 \leq i, j, k, l \leq m$. We can prove the follwoing theorem.

Theorem 20 ([5])**.** *For all $\sigma \in S_m$, we have*

$$\begin{aligned} \det\left(E + \natural_m - zI_m\right) &= \mathrm{rdet}(E + \natural^* - zI_m) \\ &= \mathrm{Det}(E + \natural_\sigma - (z+1)I_m). \end{aligned}$$

The above has the following implication.

Corollary 21. *Let $\overline{C}^{\varphi}(z)$ be the Capelli element for φ of the group algebra. For all $\sigma \in S_m$, we have*

$$\begin{aligned} \overline{C}^{\varphi}(z) &= \mathrm{rdet}(E^{\varphi} + \alpha_m\natural^* - zI_m) \\ &= \mathrm{Det}(E^{\varphi} + \alpha_m\natural_\sigma - (z+1)I_m). \end{aligned}$$

Acknowledgments I am deeply grateful to Prof. Hiroyuki Ochiai, Prof. Tôru Umeda and Prof. Minoru Itoh who provided the helpful comments and suggestions. Also, I would like to thank my colleagues in the Graduate School of Mathematics of Kyushu University, in particular Yuka Suzuki for comments and suggestions. This work was supported by a grant from the Japan Society for the Promotion of Science (JSPS KAKENHI Grant Number 15J06842).

REFERENCES

[1] ARTIN, Emil. *Geometric algebra*. Courier Dover Publications, 2016.
[2] ASLAKSEN, Helmer. Quaternionic determinants. *The Mathematical Intelligencer*, 1996, 18.3: 57–65.
[3] HUANG, An. Noncommutative multiplicative norm identities for the quaternions and the octonions. *arXiv preprint arXiv:1102.2657*, 2011.
[4] STEINBERG, Benjamin. *Representation theory of finite groups: an introductory approach*. Springer Science & Business Media, 2011.
[5] ITOH, Minoru; UMEDA, Tôru. On central elements in the universal enveloping algebras of the orthogonal Lie algebras. Compositio Mathematica, 2001, 127.03: 333–359.
[6] UMEDA, Tôru. On the proof of the Capelli identities. *Funkcialaj Ekvacioj*, 2008, 51.1: 1–15.
[7] UMEDA, Tôru. Remarks on the Capelli identities for reducible modules, preprint(2016).

Naoya Yamaguchi
Graduate School of Mathematics
Kyushu University
Nishi-ku, Fukuoka 819-0395
Japan
n-yamaguchi@math.kyushu-u.ac.jp